

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(51) International Patent Classification<sup>6</sup>:

C12Q 1/66

A1

(11) I

WO 9607759A1

(43) International Publication Date: 14 March 1996 (14.03.96)

(21) International Application Number: PCT/GB95/02085

(22) International Filing Date: 4 September 1995 (04.09.95)

(30) Priority Data:

9417793.8

5 September 1994 (05.09.94)

GB

(71) Applicant (for all designated States except US): CELSIS  
INTERNATIONAL PLC [GB/GB]; Science Park, Milton  
Road, Cambridge CB4 4FX (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FOOTE, Nicholas, Peter,  
Martin [GB/GB]; 15 The Woodlands, Linton, Cambridge  
CB1 6UF (GB). NOBLE, Michael [GB/GB]; 41 River  
Lane, Cambridge CB5 8HP (GB). GRANT, Peter, Leonard  
[GB/GB]; 28 Russett Avenue, Needingworth, Cambs PE17  
3UE (GB). EVANS, Christopher, Thomas [GB/GB]; Stable  
Cottage, Fowlmere Road, Heydon, Hertfordshire SG8 8PU  
(GB).(74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7  
Eldon Street, London EC2M 7LH (GB).(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,  
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE,  
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN,  
MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT,  
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,  
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,  
ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD,  
SZ, UG).

## Published

*With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*

(54) Title: BIOLUMINESCENCE AND ASSAY REAGENTS

## (57) Abstract

A kit of two more containers, suitable for use in the estimation of ATP released from microorganisms, contains luciferin, luciferase, a reconstitution medium and, if desired, a surfactant or other cell lytic agent, and additionally contains a medium-chain alkanolic acid.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

### BIOLUMINESCENCE AND ASSAY REAGENTS

This invention relates to a method for detecting microorganisms.

Live microorganisms can be detected with great  
5 sensitivity by measuring their ATP using a reagent based on  
the firefly luciferase reaction. This ATP must be  
extracted by lysing the microorganisms. There are  
presently four main ways of doing this, i.e. extracting  
with boiling buffer or with detergent, or using solvents or  
10 acids.

The detergent method is understandably the most  
popular because it involves a minimum of sample handling,  
is almost instantaneous, and allows a simple two-step  
assay, comprising: (1) adding an extractant to the sample;  
15 and (2) adding reagent and measuring light. This method is  
the subject of GB-A-1604249.

The detergent often actually stimulates the initial  
light production, but also rapidly inactivates the enzyme.  
Thus the light output falls to nothing over a period of a  
20 few minutes. Therefore, despite the initial stimulation,  
there may be a drastic loss of assay sensitivity,  
especially if the light output is not measured straight  
away.

It is known that non-ionic detergents protect, at  
25 least in part, from this inactivation. However, they may  
also protect bacteria from the lytic effects of the  
cationic detergent; therefore, a "mixed" detergent  
extractant is not efficient. Furthermore, non-ionic  
detergents themselves have a slow inactivating effect on  
30 luciferase and cannot be included as part of a stable  
reagent.

The only way to use a non-ionic detergent as a  
protectant is to add it between the additions of extractant  
and reagent. The assay now requires three steps, and is  
35 therefore less convenient and more expensive to automate.

An object behind the present invention is to provide  
a reagent having a low decay rate, i.e. <5% per minute,

e.g. even after the addition of a detergent-based extractant, so that it can be used in a two-step assay. This could be used in even the simplest luminometer, e.g. a camera-type microtitre plate system, and would bring all  
5 the advantages of improved accuracy and sensitivity of a true low-decay rate assay.

Surprisingly, it has been found that medium chain-length fatty acids, e.g. C<sub>5-15</sub> alkanolic acids, are efficient luciferase protectants. Their esters or other  
10 derivatives/analogues may also be used. For example, a reagent containing octanoic acid retains low-decay rate kinetics even in the presence of an efficient extractant based on cationic detergents. It has little effect on the storage stability of luciferase and can even be added to a  
15 luciferase-luciferin reagent before it is freeze-dried, although the, say, octanoic acid is preferably included in the reagent reconstitution buffer. Such a reagent could still be used after 8 hours at 25°C or 1 week at 4°C.

The following Examples illustrate the invention.

20 Example 1

Vials of a commercially-available freeze-dried reagent preparation containing firefly luciferase, D-luciferin and stabilisers, a low decay rate cocktail, were dissolved in 5 ml of buffer without and with added octanoic acid. The  
25 latter comprised 50 mM Tris-Hepes, 9 mM magnesium acetate, 1 mM EDTA, 1 g/l octanoic acid and 0.02% sodium azide. After incubation at 15°C for 1 hour, the two preparations were compared in an assay of ATP in the presence of a bacterial lysing reagent containing cationic detergents.

30 The assay was performed as follows using a Berthold Biolumat LB9500T luminometer: 0.1 ml of the lysing agent was placed in a cuvette, 0.1 ml of  $2 \times 10^{-9}$  M Na<sub>2</sub>ATP in sterile water was added, and the cuvette was equilibrated at 25°C for 5 minutes. 0.1 ml of the reagent was then  
35 added. The initial light output was measured by a 10 second integration, and the light signal was continuously monitored on a chart recorded so that the decay rate could

be calculated. The stability of the reconstituted reagents was compared by storage at 25°C and 4°C. The results were as follows:

5		Without octanoic acid	With octanoic acid
	Response to ATP (RLU/attomol)	0.62	0.42
	Decay rate (% min <sup>-1</sup> )	27	<3
	Stability 24 hr 25°C (%)	87	60
10	Stability 1 week 4°C (%)	84	76

These results show that the presence of octanoic acid provides adequate stability in solution for most applications.

15 Example 2

Vials of a high sensitivity cocktail were reconstituted and assayed as above, except that the concentration of the Na<sub>2</sub>ATP solution was  $2 \times 10^{-10}$  M. The decay of the light signal was slower in the presence of octanoic acid, and displayed single exponential kinetics. By determining the half-life of decay and multiplying by the initial light output, an accurate calculation of the total light emitted by the reaction mixture (in arbitrary units) could be made. This was shown to be a more accurate measure of the quantity of ATP than the initial light output alone.

CLAIMS

1. A kit of two more containers, suitable for use in the estimation of ATP released from microorganisms, containing luciferin, luciferase, a reconstitution medium and, if  
5 desired, a surfactant or other cell lytic agent, which additionally contains a medium-chain alkanolic acid.
2. A kit according to claim 1, wherein one container contains the medium and the acid.
3. A kit according to claim 1 or claim 2, wherein the  
10 acid is octanoic acid.
4. A kit according to any preceding claim, wherein one container contains a freeze-dried preparation including luciferin and luciferase.
5. A kit according to any preceding claim, wherein the  
15 medium is a buffer having substantially the optimum pH for the luciferin-luciferase reaction.
6. Any one of the containers defined in any of claims 1 to 5 that contains the acid.
7. An assay of cells, which comprises lysing the cells  
20 and measuring the ATP released by means of bioluminescence in the presence of a medium-chain alkanolic acid.

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12Q1/66

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 33, 25 November 1993 MD US, pages 24734-24741, W. A. FRANCISCO ET AL. 'Interaction of bacterial luciferase with aldehyde substrates and inhibitors.' see the whole document --- -/--	1-7

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \* "A" document defining the general state of the art which is not considered to be of particular relevance
- \* "E" earlier document but published on or after the international filing date
- \* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \* "O" document referring to an oral disclosure, use, exhibition or other means
- \* "P" document published prior to the international filing date but later than the priority date claimed

- \* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* "&" document member of the same patent family

Date of the actual completion of the international search

9 January 1996

Date of mailing of the international search report

16.01.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Griffith, G



C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 110, no. 15, 10 April 1989 Columbus, Ohio, US; abstract no. 131899, D. M. BYERS. 'Elongation of exogenous fatty acids by the bioluminescent bacterium Vibrio harveyi.' page 402; column 1; see abstract & J. BACTERIOL., vol. 171, no. 1, pages 59-64, ----	1-7
A	EP,A,0 309 429 (LIFE SCIENCE INTERNATIONAL AB.) 29 March 1989 see the whole document ----	1-7
A	FR,A,2 699 930 (P. KABORE) 1 July 1994 ----	
A	US,A,3 666 631 (E. RICH ET AL.) 30 May 1972 ----	
A	US,A,3 423 290 (R. C. SEAMANS, JR. ET AL.) 21 January 1969 -----	

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-309429	29-03-89	AU-B- 2484988 SE-A- 8703675 WO-A- 8902929	18-04-89 24-03-89 06-04-89
FR-A-2699930	01-07-94	NONE	
US-A-3666631	30-05-72	NONE	
US-A-3423290	21-01-69	NONE	